

U. S. AIR FORCE  
PROJECT RAND  
RESEARCH MEMORANDUM

SOLUTION SETS FOR GAMES ON THE SQUARE

I. Glicksberg and O. Gross

RM-901

8 August 1952


Assigned to \_\_\_\_\_

This is a working paper. It may be expanded,  
modified, or withdrawn at any time.

---

*The* **RAND** *Corporation*  
1700 MAIN ST. • SANTA MONICA • CALIFORNIA

Copyright 1952  
The RAND Corporation



Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>08 AUG 1952</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1952 to 00-00-1952</b>	
4. TITLE AND SUBTITLE <b>Solution Sets for Games on the Square</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Rand Corporation, Project Air Force, 1776 Main Street, PO Box 2138, Santa Monica, CA, 90407-2138</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>15</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			



Summary: Some necessary and sufficient\* conditions that a pair of non-void weak closed convex sets of strategies form the solution set of a game with continuous payoff on the square are given.

## SOLUTION SETS FOR GAMES ON THE SQUARE

I. Glilksberg and O. Gross

Let  $K$  denote the set of all optimal strategies for one player,  $L$  the corresponding set for his opponent in a game. We shall refer to  $K \times L$ , the set of all pairs  $(f, g)$ ,  $f \in K$ ,  $g \in L$ , as the solution set of the game. Any non-void weak\* closed convex set  $K$  is the set of all optimal strategies for one player in some game with continuous payoff, as was shown in [1], but of course not all pairs  $K, L$  of such sets will yield solution sets. By means of constructions similar to those used in [1] we shall determine which pairs do occur in terms of the spectra,<sup>1)</sup>  $\sigma K, \sigma L$  of these sets and the number of independent containing hyperplanes.

1. Preliminaries. As was shown in [1], any non-void weak\* ( $w^*$ ) closed convex set  $K$  of strategies is the intersection

---

1)  $\sigma K = \bigcup_{f \in K} \sigma(f)$ , which is easily seen to be a closed set.

of a sequence of half spaces, which we may express by

$$(1) \quad K = \{f \mid (\varphi_n, f) = \int \varphi_n(x) df(x) \geq 0, n = 1, \dots\}$$

where  $\{\varphi_n\}$  is a sequence of continuous functions and we may assume, for each  $n$ ,  $(\varphi_n, f) = 0$  for some  $f$  in  $K$ . Certain of these  $\varphi$ 's will yield  $(\varphi, f) = 0$  for all  $f$  in  $K$ , and these we shall denote by  $p$ 's. Thus we shall write

$$K = S(\varphi_m; p_n)^2)$$

to express the fact that  $K = \{f \mid (\varphi_m, f) \geq 0 = (p_n, f)\}$  as well as the fact that  $(\varphi_m, K)$  is a non-degenerate interval. The functions  $p_m$  thus define hyperplanes containing  $K$  while the  $\varphi_m$  do not. If the set  $K$  is the intersection of a set of hyperplanes, one may show exactly as in the proof of (1) that it is the intersection of a sequence of these and one may write  $K = S(p_m)$ .

What we shall be concerned with in large part in the following constructions will be the hyperplanes containing  $K$ . It is immediately evident that if we select from the functions  $\{p_n\}$  a maximal subsequence  $\{p'_n\}$  which is linearly independent on  $\sigma K$  then the relations  $(p_n, f) = 0$  are consequences of the relations  $(p'_n, f) = 0$  for  $f$  for which

---

2) For the opponent we shall write  $L = S(\psi_m; q_n)$  where we take  $(\psi_m, g) \neq 0$ .

$\sigma(f) \subset \sigma K$ . Consequently if we set  $p^*(x) = \text{dist}(x, \sigma K)$ , then  $(p^*, f) = 0$ ,  $(p'_n, f) = 0$  all  $n \iff (p_n, f) = 0$  all  $n$ ; thus in most of what follows we shall assume the  $\{p_n\}$  to be linearly independent,<sup>3)</sup> and actually orthonormal:

Suppose we define a measure on  $\sigma K$  in the following way: select a sequence  $\{x_n\}$  dense in  $\sigma K$  and place weight  $2^{-n}$  at  $x_n$ . Then clearly we may apply the Gram-Schmidt process to the  $\{p_n\}$  to obtain an orthonormal sequence  $\{p'_n\}$  of the same length (we take  $\{x_n\}$  dense to insure that only the function 0 has the integral of its square zero). Just as clear is the fact that  $(p_n, f) = 0$  for all  $n$  is equivalent to  $(p'_n, f) = 0$  for all  $n$ .

2. Constructions. We shall now construct payoffs which will have three types of solution sets. That these are the only types which occur will be shown later.

Case I: Suppose  $\sigma K = [0, 1] = \sigma L$  and  $K$  and  $L$  are the intersections of the same number of independent hyperplanes. The orthonormal sequences  $\{p_n\}$  and  $\{q_n\}$  defining  $K$  and  $L$  are thus of the same length, and if we set

$$M(x, y) = \sum a_n p_n(x) q_n(y),$$

---

3) We shall say that the hyperplanes  $H_n$  defined by  $H_n = \{f \mid (p_n, f) = 0\}$  are independent hyperplanes containing  $K$  if the  $p_n$  are linearly independent on  $\sigma K$ ,  $K \subset H_n$ .

where the  $a_n$  are chosen to insure uniform convergence of the series, then for  $f$  in  $K$  and  $g$  in  $L$ ,

$$\int Mdf = \sum a_n (p_n, f) q_n(y) = 0 = \sum a_n p_n(x) (q_n, g) = \int Mdg;$$

on the other hand, if  $f$  is optimal

$$\int Mdf = \sum a_n (p_n, f) q_n(y) = 0,$$

and in view of the orthogonality of the  $q_n$ ,  $f$  is in  $K$ .

Similarly every optimal  $g$  is in  $L$ , and  $K \times L$  is the solution set.

Case II: Suppose  $\sigma K = [0, 1] \neq \sigma L$ , and  $K = S(\phi_m; p_n)$ ,  $L = S(q_n)$  where there are at least as many independent hyperplanes containing  $K$  as there are containing  $L$  (thus we may assume that a maximal linearly independent set of  $p_n$ 's is at least as long as the set of  $q_n$ 's linearly independent on  $\sigma L$ ). Since  $\sigma L$  is not the full unit interval we may select an open interval  $I$  which has one end point  $y_0$  in  $\sigma L$ . Select a disjoint sequence  $\{I_n\}$  of open subintervals of  $I$  for which  $\text{dist}(y_0, I_n) \rightarrow 0$ , and an open subinterval  $I_n^*$  of each  $I_n$  whose closure lies entirely in  $I_n$ . Let  $k_n$  be a continuous non-negative function which vanishes outside  $I_n$  but is non-zero inside  $I_n$ , and which assumes the value 1 at a point  $y_n$  of  $I_n^*$ . Define a continuous function  $m_n$  which vanishes at  $y_n$  and outside  $I_n^*$ , but takes on the values  $\pm 1$ .

If we then set  $q(y) = \text{dist}(y, \sigma L \cup I_n^*)$ , then for every  $y$  not in  $\sigma L$  one of the non-negative functions  $q, k_n$  is non-zero at  $y$ .

We now define our payoff as follows: we divide the sequence  $\{p_n\}$  into  $\{p_n\}$ , orthonormal and of the same length as the  $\{q_n\}$ , and  $\{p'_n\}$ . If either of the sequences  $\{p'_n\}$  or  $\{\varphi_n\}$  are finite we use repetitions to form a sequence, and if there are no  $\varphi_n$ 's say, we take  $\varphi_n = 1$  for all  $n$ . We set (for  $b_n > 0$ , chosen to insure uniform convergence)

$$M(x, y) = \sum a_n p_n(x) q_n(y) + \sum b_n [k_n(y) \varphi_{N_n}(x) + n m_n(y) p'_{N_n}(x)] + q(y)$$

where  $\{N_n\}$  is an enumeration of the integers in which each integer occurs infinitely often. For  $f$  in  $K$  and  $g$  in  $L$

$$\int Mdf = \sum b_n k_n(y) (\varphi_{N_n}, f) + q(y) \geq 0 = \int Mdg,$$

so that both are optimal.

Suppose  $f$  is optimal; then for  $y$  in  $\sigma L$ ,

$$\sum a_n (p_n, f) q_n(y) = 0$$

whence  $(p_n, f) = 0$ , and thus

$$0 \leq \sum b_n [k_n(y) (\varphi_{N_n}, f) + n m_n(y) (p'_{N_n}, f)] + q(y),$$

and at setting  $y = y_n$ ,  $b_n (\varphi_{N_n}, f) \geq 0$ , so that  $(\varphi_n, f) \geq 0$  for all  $n$ . For  $y$  in  $I_n^*$  we have

$$0 \leq b_n [k_n(y) (\varphi_{N_n}, f) + n m_n(y) (p'_{N_n}, f)]$$



whence  $0 \leq (\varphi_{N_n}, f) + m_n(y)(p'_{N_n}, f)$ , and since  $m_n$  assumes the values  $\pm 1$ ,

$$(\varphi_{N_n}, f) \geq \pm n(p'_{N_n}, f),$$

hence

$$(\varphi_{N_n}, f) \geq n|(p'_{N_n}, f)|.$$

Since  $N_n$  takes on the value  $n_0$  infinitely often,  $(\varphi_{n_0}, f) \geq n |(p'_{n_0}, f)|$  for arbitrarily large  $n$ , and  $(p'_{n_0}, f) = 0$  for each  $n_0$ . Thus  $f$  is in  $K$ .

If  $g$  is optimal, then for any  $f$  in  $K$ ,

$$0 = \iint Mdfdg = \sum b_n(k_n, g)(\varphi_{N_n}, f) + (q, g).$$

But each term of this sum is non-negative ( $(k_n, g) \geq 0$  since  $k_n \geq 0$ ) so that surely  $(q, g) = 0$ . If  $(k_n, g) > 0$  for some  $n$  then since there is an  $f$  in  $K$  for which  $(\varphi_n, f) > 0$ , we would have a contradiction. Thus

$$(q, g) = 0, (k_n, g) = 0, \text{ and } (m_n, g) = 0$$

since  $(k_n, g) = 0$  implies  $g$  places no weight on  $I_n$ . Thus  $\sigma(g) \subset \sigma L$ , and since we now may write

$$0 = \sum a_n p_n(x)(q_n, g), \quad x \text{ in } \sigma K,$$

and  $(q_n, g) = 0$ ,  $g$  is in  $L$ .

Case III:  $\sigma K \neq [0,1] \neq \sigma L$ . Here we may take any  $K$  and  $L$  without further restriction, so that  $K = S(\psi_m; p_n)$  and  $L = S(\psi_m; q_n)$  ( $(\psi_m, g) \leq 0$  here, however, in our definitions). We construct functions  $h_n$  similar to the  $k_n$  of case II, and  $\ell_n$  similar to the  $m_n$ , on an interval abutting  $\sigma K$ . We set

$$M(x, y) = \sum a_n [h_n(x) \psi_{N_n}(y) + n \ell_n(x) q_{N_n}(y) + k_n(y) \psi_{N_n}(x) + n m_n(y) p_{N_n}(x)] .$$

Arguments entirely similar to those used in case II show  $K \times L$  to be the solution set.

3. Generality. In case I ( $\sigma K = [0,1] = \sigma L$ ) we restricted our attention to the case in which  $K$  and  $L$  were intersections of the same number of independent hyperplanes. Suppose now that a game with payoff  $M$  has as its solution set  $K \times L$  where  $\sigma K$  and  $\sigma L$  are the full intervals.  $K$  is determined as the set of all  $f$  for which

$$\int M(x, y) df(x) = 0$$

(for convenience we take the value to be zero), and thus is the intersection of hyperplanes given by the functions  $\{M(\cdot, y)\}$ , and similarly  $L$  is the intersection of the hyperplanes determined by the functions  $\{M(x, \cdot)\}$ .

If a maximal linearly independent set  $\{M(x_i, \cdot)\}$  of the first set, say, is finite,  $i = 1, \dots, n$ , then the same is true of the second, indeed there are just as many. For, as is

well known,  $n$  functions  $F_1, \dots, F_n$  are linearly independent on a set  $X$  if and only if there exist  $x_1, \dots, x_n$  in  $X$  for which

$$\det (F_i(x_j)) \neq 0;$$

consequently we have  $y_1, \dots, y_n$  for which

$$(2) \quad \det (M(x_i, y_j)) \neq 0,$$

so that the functions  $\{M(\cdot, y_j)\}_{j=1, \dots, n}$  are linearly independent.

Of course if  $\{M(\cdot, y_j)\}_{j=1, \dots, n+1}$  were linearly independent

by the same argument we should have an  $x_{n+1}$  for which

$$\{M(x_i, \cdot)\}_{i=1, \dots, n+1} \text{ were, which contradicts our assumption,}$$

and there are exactly  $n$ . Thus the type of solution sets

considered in case I are the only type which can occur. (One

might note that here finite set of independent containing

hyperplanes can only occur in a polynomial-like game, since for

every  $x$  we have coefficients  $a_i(x)$  for which

$$M(x, y) = \sum a_i(x) M(x_i, y),$$

and (2) shows the functions  $a_i$  to be continuous.)

In case II,  $(\sigma K = [0, 1] \neq \sigma L)$  we considered only those  $K$  and  $L$  for which we had as many independent hyperplanes containing  $K$  as there are containing  $L$ . But if  $M$  is the payoff of a game with solution set  $K \times L$ ,  $\sigma K = [0, 1] \neq \sigma L$ , then as before since  $L$  is determined by

$$\int M(x, y) dg(y) = 0, \quad \text{all } x,$$

$L$  is just the intersection of hyperplanes. If there are only  $n$  independent containing hyperplanes, then, as we shall see in a moment, these must be given by the functions

$\{M(x_i, \cdot)\}_{i=1, \dots, n}$  linearly independent on  $\sigma L$ , for some set  $x_1, \dots, x_n$ ; consequently there exist  $y_1, \dots, y_n$  in  $\sigma L$  for which (2) holds, and  $\{M(\cdot, y_j)\}_{j=1, \dots, n}$  are linearly independent.

Since  $\int M(x, y) df(x) = 0$  for  $y$  in  $\sigma L$ , these functions define  $n$  independent hyperplanes containing  $K$ .

To see that the  $n$  independent hyperplanes containing  $L$  arise from functions  $M(x_i, \cdot)$  we note that for each  $x$ ,  $M(x, \cdot)$  defines a containing hyperplane since  $x$  is in  $\sigma K = [0, 1]$ . Consequently there can be only  $m$  points,  $m \leq n$ ,  $x_1, \dots, x_m$  for which  $\{M(x_i, \cdot)\}$  are linearly independent, so that clearly  $L = \{g | (M(x_i, \cdot), g) = 0, i = 1, \dots, m\}$ .

If  $m < n$ , we can find a function  $q_0$  for which, denoting  $M(x_i, \cdot)$  by  $q_i$ , the set  $q_0, \dots, q_m$  is linearly independent on  $\sigma L$  and  $(q_0, g) = 0$  for all  $g$  in  $L$ . But then the mapping

$$T: g \rightarrow ((q_0, g), \dots, (q_m, g)),$$

of the set  $S$  of all strategies into  $m + 1$  space, takes  $S$  into a convex subset containing  $(0, \dots, 0)$  (since  $L$  is non-void).

But  $T(S)$  intersects the line  $(t, 0, \dots, 0)$  in only one point (since  $(q_0, g) = 0$  for  $g$  in  $L$ )—thus  $(0, 0, \dots, 0)$  is a boundary point and we have a supporting hyperplane at this point given by constants (not all zero)  $a_0, \dots, a_m$ . Thus  $\sum_{i=0}^m a_i (q_i, g) \geq 0$

for all  $g$  in  $S$ , hence  $\sum a_i q_i(y) \geq 0$  for  $y$  in  $\sigma L$ . If inequality holds for any  $y$  it holds in some neighborhood, and this is, of course, of positive measure with respect to some  $g$  in  $L$  (from the definition of  $\sigma L$ ), whence  $\sum a_i(q_i, g) > 0$  for some  $g$  in  $L$  - a contradiction. Thus  $\sum a_i q_i = 0$  on  $\sigma L$ , which contradicts the linear independence on  $\sigma L$ , and we must have  $m = n$ .

Thus the theme of things is as follows: The necessary and sufficient condition that  $K \times L$  be the solution set for a game with continuous payoff on the square (where  $K$  and  $L$  are non-void  $\omega^*$  closed convex sets of strategies) is that one of the following hold:

- (a)  $\sigma K = [0, 1] = \sigma L$  and  $K$  and  $L$  are the intersection of the same number (finite if and only if the game is polynomial-like) of independent containing hyperplanes
- (b)  $\sigma K = [0, 1] \neq \sigma L$ ,  $L$  is the intersection of hyperplanes and  $K$  has as many independent containing hyperplanes as  $L$
- (c)  $\sigma K \neq [0, 1] \neq \sigma L$ .

The constructions we have used can be duplicated in the case of a game with continuous payoff played on a pair of infinite compact metric spaces; the character of solution sets, however, involves slightly different conditions:

$\sigma K = [0,1] = \sigma L$  must be replaced by  $\sigma K$ ,  $\sigma L$  open,  
 $\sigma K = [0,1] \neq \sigma L$  by  $\sigma K$  open,  $\sigma L$  not open,  
 $\sigma K \neq [0,1] \neq \sigma L$  by  $\sigma K$  and  $\sigma L$  not open. In the case of  
 a unique optimal strategy forming  $K$  and another forming  $L$   
 we are thus guaranteed a game having  $K \times L$  as the solution  
 set, which generalizes the result of [2].

As a final remark, we note that solution sets for  
 symmetric games on the square (where  $M(x,y) = -M(y,x)$ ) can be  
 easily described. For such games the value is always zero  
 and any optimal strategy for one player is optimal for his  
 opponent, so that a solution set is of the form  $K \times K$ . The  
 necessary and sufficient condition that  $K \times K$  be the solution  
 set of a symmetric game is that either

- (a)  $\sigma K = [0,1]$  and  $K$  is the intersection of an even  
 (we take  $\infty$  as even) number of independent hyper-  
 planes, or
- (b)  $\sigma K \neq [0,1]$ .

For if  $\sigma K = [0,1]$  and  $K$  is the intersection of an  
 even number of independent hyperplanes given by functions  
 $\{p_n\}$  (which we may take orthonormal), then, dividing these  
 into two sets  $\{p_n\}$ ,  $\{p'_n\}$  of equal cardinality, we may set

$$M(x,y) = \sum a_n [p_n(x)p'_n(y) - p'_n(x)p_n(y)],$$

which is easily seen to have  $K \times K$  as its solution, and is  
 symmetric. On the other hand, if  $K \times K$  is the solution set

of a game with payoff  $M$  and  $\sigma K = [0,1]$ , then  $K$  is, of course, the intersection of a set of hyperplanes. If only a finite number of these are independent, then, as before,  $M$  is polynomial-like, that is,

$$M(x,y) = \sum_{n=1}^k \varphi_n(x) \psi_n(y),$$

where  $\{\varphi_n\}$  and  $\{\psi_n\}$  are linearly independent sets of functions. Since  $M$  is symmetric

$$M(x,y) = -M(y,x) = -\sum_{n=1}^k \varphi_n(y) \psi_n(x),$$

so

$$M(x,y) = \frac{1}{2} \sum_{n=1}^k [\varphi_n(x) \psi_n(y) - \varphi_n(y) \psi_n(x)].$$

If the functions  $\{\varphi_n, \psi_n\}$  are not a linearly independent set, replacement of a dependent  $\varphi$  or  $\psi$  again yields a sum of the same type, and we finally obtain a similar expression for  $M$  in which the set  $\{\varphi_n, \psi_n\}$  is linearly independent; however, there are an even number of terms in the resulting sums, and thus there must be an even number of independent hyperplanes determining  $K$ .

In case (b),  $K = S(\varphi_m; p_n)$ , and we may set

$$M(x,y) = \sum a_n [k_n(y)\varphi_{N_n}(x) + n m_n(y)p_{N_n}(x) - k_n(x)\varphi_{N_n}(y) - n m_n(x)p_{N_n}(y)]$$

to obtain a symmetric game in which  $K \times K$  is the solution set.

## REFERENCES

1. I. Glicksberg and O. Gross, Optimal Sets for Games over the Square, RM-889.
2. I. Glicksberg and O. Gross, Continuous Games with Given Unique Solutions, RM-620.